



National Day Nurseries Association

**★Brighter thinking
for early years**

Quiz

A young girl with light hair and blue eyes is shown in profile, looking towards the left. She is wearing a red scarf and a white sweater. A stethoscope is around her neck, and she is holding the earpieces to her ears. To her left is a large, brown teddy bear. The background is a soft-focus outdoor scene with yellow autumn leaves. The bottom half of the image is overlaid with a dark blue circular graphic.

Paediatric First Aid



National Day Nurseries Association

***Brighter thinking
for early years**

NDNA Quiz: Paediatric first aid (NDNA and Millie's Mark)

Question 1: Can an AED (automated external defibrillator) be used on a baby or child?

- Yes, but only if it has 'paediatric' pads or a paediatric setting
- No, it should never be used
- Yes, and you can use adult pads

Question 2: How would you recognise a broken or dislocated bone?

- Pain, loss of power and unnatural movement
- Swelling/bruising, deformity and crepitus (joint sounds such as popping)
- All of these

Question 3: A baby is choking. You have followed the initial steps but the obstruction has not cleared – how do you give chest thrusts?

- Do not give chest thrusts to a baby – it is too dangerous
- Use one finger to give up to ten chest thrusts
- Use two fingers to give up to five chest thrusts

Question 4: The baby has become unresponsive and isn't breathing normally. You have asked someone to call 999/112 and are about to give rescue breaths and chest compressions. How many breaths and compressions? Give five initial rescue breaths and use two fingers to give 30 compressions

- Give ten initial breaths and use two fingers to give 50 compressions
- Give five initial rescue breaths and use three fingers to give 20 compressions

Question 5: What should you do if a nosebleed is severe or lasts more than 30 minutes?

- Make sure they are sitting upright and wait for another 30 minutes
- Nip the soft part of the nose
- Take the child to hospital sitting upright

Question 6: How would you treat a bruise?

- Place an ice pack straight into the bruise and apply pressure for 10 minutes
- Place an ice pack in a tea towel / triangular bandage on the bruise and apply pressure for ten minutes
- Place an ice pack on the bruise and ask the child to hold it there

Question 7: What are the 'red flag' early warning signs of meningitis?

- Cold hands and feet, abnormal skin colouring, pain in the limbs and joints
- Fever, blotchy purple rash, drowsiness, headache, stiff neck and dislike of bright lights
- All of the above

Question 8: How urgently should you seek medical advice if you suspect a child has meningitis?

- Wait for symptoms to appear
- Seek urgent medical advice
- Wait for parents to arrive

Answers – yellow marks the correct answer (s)

Question 1: Can an AED (automated external defibrillator) be used on a baby or child?

- Yes, but only if it has 'paediatric' pads or a paediatric setting
- No, it should never be used
- Yes, and you can use adult pads

Note: If you only have adult pads, you should use them as this is better than not using an AED at all in an emergency.

Question 2: How would you recognise a broken or dislocated bone?

- Pain, loss of power and unnatural movement
- Swelling/bruising, deformity and crepitus (joint sounds such as popping)
- All of these

Question 3: A baby is choking. You have followed the initial steps but the obstruction has not cleared – how do you give chest thrusts?

- Do not give chest thrusts to a baby – it is too dangerous
- Use one finger to give up to ten chest thrusts
- Use two fingers to give up to five chest thrusts

Note: If the obstruction does not clear, repeat the back slaps and chest thrusts and if it is ineffective then should for help and have someone call 999/112.

Question 4: The baby has become unresponsive and isn't breathing normally. You have asked someone to call 999/112 and are about to give rescue breaths and chest compressions. How many breaths and compressions?

- Give five initial rescue breaths and use two fingers to give 30 compressions
- Give ten initial breaths and use two fingers to give 50 compressions
- Give five initial rescue breaths and use three fingers to give 20 compressions

Question 5: What should you do if a nosebleed is severe or lasts more than 30 minutes?

- Make sure they are sitting upright and wait for another 30 minutes
- Nip the soft part of the nose
- Take the child to hospital sitting upright

Question 6: How would you treat a bruise?

- Place an ice pack straight into the bruise and apply pressure for 10 minutes
- Place an ice pack in a tea towel / triangular bandage on the bruise and apply pressure for ten minutes
- Place an ice pack on the bruise and ask the child to hold it there

Question 7: What are the 'red flag' early warning signs of meningitis?

- Cold hands and feet, abnormal skin colouring, pain in the limbs and joints
- Fever, blotchy purple rash, drowsiness, headache, stiff neck and dislike of bright lights
- All of the above

Note: Early signs: Cold hands and feet, abnormal skin colouring, pain in the limbs and joints

Later signs: Fever, blotchy purple rash, drowsiness, headache, stiff neck and dislike of bright lights

Question 8: How urgently should you seek medical advice if you suspect a child has meningitis?

- Wait for symptoms to appear
- Seek urgent medical advice
- Wait for parents to arrive

Note: Time is critical – do not wait for symptoms to appear. Call 999/112 for emergency help if a rash is present and go straight to hospital if no medical help is available.

To learn more, visit our Millie's Mark webpage: <https://www.milliesmark.com/>

The contents of this work are exclusively the property of National Day Nurseries Association (NDNA). We strive to ensure that this content is accurate and up-to-date, but information can change over time, so this document should not be used as a substitute for personalised advice from a qualified professional. We hope that you will continue to use this quiz as a useful tool and would welcome any feedback. If you have any queries or concerns about the quiz please do not hesitate to contact us.

National Day Nurseries Association

National Early Years Enterprise Centre, Longbow Close, Huddersfield HD2 1GQ
tel: 01484 407070 fax: 01484 407060 info@ndna.org.uk www.ndna.org.uk